arXiv:1211.5597v1 [quant-ph] 22 Nov 2012

Effects on the Non-Relativistic Dynamics of a Charged Particle Interacting with a Chern-Simons Potential

F. Caruso[a,b], J.A. Helayël-Neto[a], J. Martins[b], V. Oguri[b]

[a]*Centro Brasileiro de Pesquisas Físicas - Rua Dr. Xavier Sigaud, 150, 22290-180, Urca, Rio de Janeiro, RJ, Brazil*
[b]*Instituto de Física Armando Dias Tavares, Universidade do Estado do Rio de Janeiro - Rua São Francisco Xavier, 524, 20550-900, Maracanã, Rio de Janeiro, RJ, Brazil*

Abstract

The hydrogen atom in two dimensions, described by a Schrödinger equation with a Chern-Simons potential, is numerically solved. Both its wave functions and eigenvalues were determined for small values of the principal quantum number n The only possible states correspond to $l = 0$. How the result depends on the topological mass of the photon is also discussed. In the case $n = 1$, the energy of the fundamental state corresponding to different choice for the photon mass scale are found to be comprehended in the interval $-3,5\times10^{-3}$ eV $\leq E \leq -9,0\times10^{-2}$ eV, corresponding to a mean radius of the electron in the range $(5.637\pm0.005)\times10^{-8} \leq \langle r\rangle \leq (48.87\pm0.03)\times10^{-8}$ cm. In any case, the planar atom is found to be very weekly bounded showing some features similar to the Rydberg atoms in three dimensions with a Coulombian interaction.

Keywords: Chern-Simons potential, Schrödinger equation, hydrogen atom

---

1. Introduction

Lower-dimensional quantum-mechanical and field-theoretic models have acquired a renewed interest and have boosted a tremendous deal of papers over the few recent years, especially in connection with Condensed Matter Systems. Relevant phenomena such as high-T_c superconductivity [1, 3], fractional quantum Hall effect [4, 7] and spin-charge splitting [8, 11] in planar

---

Email address: `francisco.caruso@gmail.com` (F. Caruso)

systems have motivated an intensive development of models based on principles, fundamental ideas and technicalities of Quantum Field Theory [12]. New materials like the genuinely planar graphene monolayers [13, 14], topological insulators [15]-[19] and topological superconductors [20]-[22] bring to light what we may say is a new state of matter, the so-called topological matter. Peculiar properties, which are typical of lower-dimensional systems, are revealed in these systems which motivate the real need to apply ordinary and relativistic Quantum Mechanics, methods of Quantum Field Theory and concepts of gauge theories to build up consistent models that correctly describe the observed phenomena.

In the framework of lower-dimensional Physics, more specifically, planar systems, electromagnetic interaction allows that the photon acquires a non-trivial mass without any conflict with its inherent gauge symmetry. There emerges then the so-called Maxwell-Chern-Simons Electrodynamics [23, 24], which, as quantum gauge-field theories, display a remarkable ultraviolet behavior and finds applications in the understanding and description of a great diversity of phenomena which are typical of (1+2) space-time dimensions, such as, for the example, the appearance of particles, the anyons [25, 26], which obey an intermediate statistics and are present in many planar systems of interest.

Motivated by the importance and particular aspects of Planar Electrodynamics, we place our question of interest in this frame: namely, we wish to exploit and discuss the influence of space-time dimensions on the electromagnetic interactions and on the properties of particles with electric charge and/or magnetic properties. With this purpose in mind, our paper sets out to investigate the non-relativistic quantum mechanics of an electron in the field of a static point charge considering that the electromagnetic phenomenon is mediated by the (gauge-invariant) massive photon of a Maxwell-Chern-Simons Electrodynamics. By following this path, out true aim is to reassess properties of the non-relativistic hydrogen atom and, so, to compare them with the corresponding properties in other dimensions to really point out the effects of dimensionality on the structure of the hydrogen atom.

The outline of our paper is as follows: in Section 2, we set up the non-relativistic planar Schrödinger equation for an electron in a Maxwell-Chern-Simons potential. Next, in Section 3, we carry out the numerical studies of the problem and understand that $l = 0$ is the only possible state for the Maxwell-Chern-Simons planar atom. In this s-wave state, we work out the values of the binding energy and mean radius of the electron orbit, investigating how those values depends on the choice of the topological mass of the photon involved in the Maxwell-Chern-Simons Electrodynamics. Finally, in Section 4, we cast our Concluding Comments.

2. Hydrogen atom for Chern-Simons potential

For the classical Maxwell-Chern-Simons Electrodynamics in $(1+2)$ dimensions, the magnetic and the electric fields of a pointlike charge e are given, in the Gaussian unit system, by

$$B = -\frac{m_\gamma ce}{2\pi\hbar}\, K_\circ(m_\gamma cr/\hbar); \qquad \vec{E} = \frac{\hbar}{m_\gamma c}\vec{\nabla}B = -\vec{\nabla}\phi$$

where $K_\circ$ is the modified Bessel function and m_γ is the photon effective topological mass, which is expected to be in the range 0.1 eV $\leq m_\gamma \leq$ 1 eV [27]. From these equations we get the electric potential

$$\phi(r) = \frac{em_\gamma c}{2\pi\hbar}\, K_\circ(m_\gamma cr/\hbar)$$

In two dimensions, the Schrödinger equation for a massive charged particle (electron), with mass m and charge e, interacting with a central external potential ϕ is

$$\frac{\mathrm{d}^2u(r)}{\mathrm{d}r^2} + \frac{2m}{\hbar^2}\left\{E - U(r) - \frac{\hbar^2}{2m}\frac{[l^2-1/4]}{r^2}\right\}u(r) = 0, \tag{1}$$

where the potential energy is $U(r) = -e\phi(r)$ and the usual radial solution (in D dimensions) is given by $R(r) = \dfrac{u(r)}{r^{(D-1)/2}}$.

Therefore, the Schrödinger equation for a Maxwell-Chern-Simons potential is

$$\frac{\mathrm{d}^2u(r)}{\mathrm{d}r^2} + \frac{2m}{\hbar^2}\left\{E + \frac{m_\gamma ce^2}{2\pi\hbar}K_0\left(\frac{m_\gamma c}{\hbar}r\right) - \frac{\hbar^2}{2m}\frac{[l^2-1/4]}{r^2}\right\}u(r) = 0. \tag{2}$$

In order to solve numerically Eq. (2), it is convenient to put it in a dimensionless form. Defining Bohr's radius as $a_B \equiv \dfrac{\hbar^2}{me^2}$, one gets

$$\frac{\mathrm{d}^2u(r)}{\mathrm{d}r^2} + \frac{1}{a_B^2}\left\{\frac{E}{e^2/2a_B} + \frac{m_\gamma ca_B}{\pi\hbar}K_0\left(\frac{m_\gamma c}{\hbar}r\right) - a_B^2\frac{[l^2-1/4]}{r^2}\right\}u(r) = 0. \tag{3}$$

Introducing the dimensionless variable $x = r/a_B$, Eq. (3) can be written as

$$\frac{\mathrm{d}^2 y(x)}{\mathrm{d}x^2} + \left\{ \frac{E}{e^2/2a_B} + \frac{m_\gamma c a_B}{\pi \hbar} K_0 \left(\frac{m_\gamma c a_B}{\hbar} x \right) - \frac{[l^2 - 1/4]}{x^2} \right\} y(x) = 0. \quad (4)$$

Finally, expressing $m_\gamma = \lambda m_e$, Eq. (4) can be also written in terms of the fine structure constant, α, as

$$\alpha = \frac{e^2}{\hbar c} = \frac{\hbar^2}{m a_B} \frac{1}{\hbar c} = \frac{\hbar}{m c a_B} = \frac{1}{137}. \quad (5)$$

from which we gets the dimensionless equation

$$\frac{\mathrm{d}^2 y(x)}{\mathrm{d}x^2} + \left\{ \frac{E}{e^2/2a_B} + \frac{\lambda}{\pi \alpha} K_0 \left(\frac{\lambda x}{\alpha} \right) - \frac{[l^2 - 1/4]}{x^2} \right\} y(x) = 0 \quad (6)$$

Thus the problem is reduced to the interaction of a charged particle with an effective potential, $U_{\text{eff}}(x)$, defined by:

$$U_{\text{eff}}(x) = -\frac{\lambda}{\pi \alpha} K_0 \left(\frac{\lambda x}{\alpha} \right) + \frac{[l^2 - 1/4]}{x^2}. \quad (7)$$

which generalizes what was done in Ref. [28].

Such an effective potential in two dimensions has a very peculiar behavior, namely, what is usually called "centrifugal potential" became indeed attractive when $l = 0$ and it is straightforward to see that only in those case it is possible to have bound states.

This effective potential, for $l = 0$, and $m_\gamma/m_e = \lambda \simeq 0.2 \times 10^{-5}$, has the behavior as shown in Fig. 1.

The eigenvalue equation, Eq. (6), can be numerically solved by a slightly modified version of the Numerov Method [30],[31], as did in Ref. [29], in order to include also first derivatives The algorithm was implemented using C^{++} language. All the calculations and graphics were done by using the CERN/ROOT package.

3. General results

First of all, it is important to say that it is not our intention to carry out a systematic calculation of the hydrogen atom sprectra; only few eigenstates



Figure 1: Plot of the effective potential and of each one of its components, for $l = 0$.

and eigenfunctions were determined. Once more, let us stress that for the Maxwell-Chern-Simons potential, the planar structure of the space imposes solutions only with $l = 0$. For different values of the angular momentum, the centrifugal potential dominates and the effective potential became positive and continuously decreases.

The second point, already mentioned, is that the result clearly depends on the choice of the topological mass of the photon, m_γ. Thus, for numerical purpose, we have considered three cases: $\lambda = m_\gamma/m_e = 0.2 \times 10^{-5}$, $\lambda = 0.2 \times 10^{-4}$ and $\lambda = 0.2 \times 10^{-3}$. For $m_\gamma \simeq 0.1$ eV we could not find a solution within the sensitivity of our numerical calculation. This situation is very close to that where only the 'centrifugal term' contributes which gives rise to no bounded state. Conversely, once some energy levels of this kind of planar atom is measured we can straightforwardly set up the scale of m_γ.

For the ground state we found numerical solutions for the all probed λ scales. The three wavefunctions are shown in Fig. 2 and the corresponding eigenvalues for each one are, respectively,

$$\begin{aligned} E_1 &= -0.000026 \text{ Ry} = -0.000026 \times (13.6) \text{ eV} = -0.0035 \text{ eV}; \\ E_1 &= -0.0015 \text{ Ry} = -0.020 \text{ eV}; \\ E_1 &= -0.0067 \text{ Ry} = -0.09 \text{ eV}. \end{aligned}$$

In the case $\lambda = 0.2 \times 10^{-5}$ ($m_\gamma \simeq 1$ eV), we still find a second state ($n = 2$, $l = 0$) with energy $E_2 = -0.0002$ Ry. For $\lambda = 0.2 \times 10^{-4}$ ($m_\gamma \simeq 10$ eV), two



Figure 2: Radial wave function $R_{l=0}(r/a_B)$ for the hydrogen atom in the case $D = 2$ with arbitrary normalization. The three curves correspond to different choice of m_γ.

other states are found ($n = 2$, $l = 0$ and n=3, $l = 0$) with energies given, respectively, by $E_2 = -0.00075$ Ry and $E_3 = -0.0004$ Ry.

Fig. 3 shows the first three eigenfunctions corresponding to $m_\gamma = 0.1$ eV.



Figure 3: Radial wave functions for the planar hydrogen atom for the states $n = 1, 2, 3$ with arbitrary normalization.

Once the radial solution is known, the mean value of the electron radius, $\langle r \rangle$, can be evaluated by solving the expression (with $D = 2$)

$$\langle r\rangle = \frac{\int r^D \cdot R^2_{l=0}(r)\mathrm{d}r}{\int r^{D-1} \cdot R^2_{l=0}(r)\mathrm{d}r} = a_B \frac{\int x \cdot y^2_{l=0}(x)\mathrm{d}x}{\int y^2_{l=0}(x)\mathrm{d}x} \tag{8}$$

which was done numerically adopting the Rejection Monte Carlo Method [33, 34].



Figure 4: Histogram generated by the Rejection Monte Carlo Method in order to estimate the mean value of the electronic radius in the ground state of the planar hydrogen atom, corresponding to the choice $m_\gamma = 1$ eV.

For the ground state, and considering $m_\gamma = 1$ eV, the result for the mean radius is $\langle x\rangle = \langle r\rangle/a_B = (92.35 \pm 0.06)$, which corresponds to the statistic treatment of the histogram given in Fig. 4. Using the value of Bohr's radius, $a_B = 0.52917720859(36) \times 10^{-8}$ cm [32] (in Gaussian unit system), the mean radius of the electron in the ground state of a planar hydrogen atom described by the Schrödinger equation with a Chern-Simons potential, is $\langle r\rangle = (48.87 \pm 0.03) \times 10^{-8}$ cm. This result is two order of magnitude greater than the atomic radius of the hydrogen atom in three dimensions ($\sim 10^{-8}$ cm). The values of the orbital radius for different choices of m_γ (λ) are shown in Table 1.

Table 1: Mean values of r/a_B by varying λ (or m_γ).

λ	$\langle r\rangle/a_B$
0.2×10^{-5}	92.35 ± 0.06
0.2×10^{-4}	25.59 ± 0.01
0.2×10^{-3}	10.653 ± 0.009

These values can be compared to the results of the same mean value for hydrogen atomic radius in higher dimensional spaces [29], from the number of dimensions varying from 3 to 10 (remember that for $D = 4$ there is no solution for the hydrogen atom) leading to the conclusion that in two dimensions the orbital radius of the electron in the ground state is by far the biggest one.

4. Final Comments

Let us compare our result with the well known features of the hydrogen atom in three dimensions. Remember that the ground state of this simple atom with such dimensionality is bound by 1 Ry, and the orbital radius of the electron is one Bohr's radius, 1 a_B. In contrast, the $n = 10$ state, which can be considered as a relatively low Rydberg state, has a binding energy of 0.01 Ry and a radius of 100 a_B. Now, if we compare our result for the ground state, bounded by the energy $E_1 = 0.00026$ Ry, with the three-dimensional analogous energy $E_n \sim 1/n^2$, it will correspond to the principal quantum number $n = 62$, in three dimensions, with a mean radius $\langle r \rangle \simeq 92 a_B$, instead of 100 a_B. Notice that the binding energy of the fundamental state of hydrogen atom in two dimensions (3.5 meV) is comparable to one tenth of thermal energies ($\simeq$ 40 meV). Therefore, one should expect that planar atoms with one electron might be useful system with which to probe the interaction of an atom and the vacuum [35]. Also the possibility of having giant molecules [36], [38], [37] could be investigated.

In two space dimensions, new aspects of the electromagnetic interactions could be considered that may enrich our study of the hydrogen atom and introduce new corrections to its binding energy and mean radius. Among these, we should consider a non-minimal coupling [39],[40],[41] through the dual of the electromagnetic field strength, along with the minimal coupling with the potential in the gauge covariant derivative. This is property specific of (1+2)-D. Also, it would be of interest, for the sake of computing effects due to the magnetic properties of the electron, to re-analyze the magnetic moment and the gyromagnetic ratio (the g-factor) of the electron in a planar scenario to get its influence on the corrections to the quantities we have numerically computed in Section 3.

It is our immediate purpose to go through this direction and we shall be soon reporting on that elsewhere.

References

[1] J.G. Berdnoz, K.A. Müller, Z. Phys. B 64 (1986) 189.

[2] P.W. Anderson, Science 235 (1987) 1196.

[3] P.W. Anderson, The Theory of Superconductivity in the High-Tc Cuprates. Princeton: University Press (1997).

[4] K. von Klitzing, G. Dorda, M. Pepper, Phys. Rev. Lett. 45 (1980) 494.

[5] R.B. Laughlin, Phys. Rev. Lett. 50 (1983) 1395.

[6] H.L. Stormer, Rev. Mod. Phys. 71 (1999) 875.

[7] R.E. Prange, S.M. Girvin, The Quantum Hall Effect, 2nd. Edition. New York: Springer-Verlag (1990).

[8] S. Tomonaga, Prog. Theor. Phys. 5 (1950) 544.

[9] J.M. Luttinger, J. Math. Phys. 4 (1963) 1154.

[10] F.D.M. Haldane, J. Phys. C: Solid State Physics 14 (1981) 2585.

[11] P.A. Marchetti, Z.B. Su, L. Yu Phys. Rev. B 58 (1998) 5808.

[12] A.M. Tsvelik, Quantum Field Theory in Condensed Matter Physics, 2nd. Edition. Cambridge: University Press (2003).

[13] A.K. Geim, K.S. Novoselov, Nature Materials 6 (2007) 183.

[14] A.K. Geim, Science 324 (2009) 1530.

[15] L. Fu, C.L. Kane, Phys. Rev. B 75 (2007) 121306 (R).

[16] H. Zhang et al., Nat. Phys. 5 (2009) 438.

[17] D. Hsieh et al., Nature (London) 452 (2008) 970.

[18] Y.L. Chen et al., Science 325 (2009) 178.

[19] X.-L. Qi, S.-C. Zhang, Rev. Mod. Phys. 83 (2011) 1057.

[20] N. Read, D. Green, Phys. Rev. B 61 (2000) 10267.

[21] X.-L. Qi, T. L. Hughes, S, Raghu, S.-C. Zhang, Phys. Rev. Lett. 102 (2009) 187001.

[22] L. Fu, E. Berg, Phys. Rev. Lett. 105 (2010) 097001.

[23] S. Deser, R. Jackiw, S. Templeton, Ann. Phys. (N.Y.) 140 (1982) 372.

[24] S. Deser, R. Jackiw, S. Templeton, Phys. Rev. Lett. 48 (1982) 975.

[25] J.M. Leinaas, J. Myrheim, Il Nuovo Cim. B 18 (1977) 1.

[26] F. Wilczek, Phys. Rev. Lett. 48 (1982) 957.

[27] T.P. Orlando, K.A. Delin, Foundations of Applied Superconductivity. Reading Massachusetts: Addison-Wesley (1991).

[28] O. Atabek, C. Deutsch, M. Lavaud, Phys. Rev. A 9 (1974) 2617.

[29] F. Caruso, J. Martins, V. Oguri, arXiv:quant-ph/1205.3740v1 (2012). Submitted for publication.

[30] B.V. Numerov, Monthly Notices of the Royal Astronomical Society 84 (1924) 592.

[31] B.V. Numerov, Astronomische Nachrichten 230 (1927) 359.

[32] Particle Data Group, Physical Review D 86 (2012) 010001.

[33] G. Cowan, Statistical Data Analysis. Oxford: Claredon Press (1998).

[34] F. James, Reports on Progress in Physics 43 (1980) 1145.

[35] T.F. Gallagher Reports on Progress in Physics 51 (1988) 143; Rydberg Atoms. Cambridge: University Press (2005).

[36] V. Bendkowsky, B. Butscher, J. Nipper, J.P. Shaffer, R. Löw, T. Pfau Nature 458 (2009) 1005.

[37] S.M. Farooqi et al. Phys. Rev. Lett. 91 (2003) 183002.

[38] C. Boisseau, I. Simbotin, E. Côté Phys. Rev. Lett. 88 (2002) 133004.

[39] S.K. Paul, A. Khare, Phys. Lett. B193 (1987) 253; erratum, *ibid.*, B196 (1987) 571.

[40] F.A.S. Nobre, C.A.S. Almeida, Phys. Lett. B455 (1999) 213.

[41] R.C. Paschoal, J.A. Helayël-Neto, Phys. Lett. A313 (2003) 412.